

G<sup>4</sup> - In an embodiment of this invention, the GABA<sub>B</sub>R1a polypeptide has an amino acid sequence identical to the amino acid sequence shown in Figures 24A-24D (SEQ ID NO: 56) and the GABA<sub>B</sub>R1b polypeptide has an amino acid sequence identical to the amino acid sequence shown in Figures 25A-25D (SEQ ID NO: 55). --

In the Claims:

Please amend claims 208, 213, 224, 230, and 231 as follows:

G<sup>5</sup> -208. (Four Times Amended) A process for determining whether a chemical compound is an agonist of a mammalian GABA<sub>B</sub>R1/R2 receptor which comprises contacting cells containing nucleic acid encoding, and expressing on their cell surface, the GABA<sub>B</sub>R1/R2 receptor, wherein such cells prior to being transfected with such nucleic acid do not express the GABA<sub>B</sub>R1/R2 receptor, with the compound under conditions permitting the activation of the GABA<sub>B</sub>R1/R2 receptor, and detecting an increase in activity of the GABA<sub>B</sub>R1/R2 receptor, wherein said increase in activity indicates that the compound is an agonist of a GABA<sub>B</sub>R1/R2 receptor, wherein the mammalian GABA<sub>B</sub>R1/R2 receptor comprises a GABA<sub>B</sub>R1 polypeptide and a GABA<sub>B</sub>R2 polypeptide, which GABA<sub>B</sub>R1 polypeptide has an amino acid sequence identical to the amino acid sequence shown in Figures 24A-24D (SEQ ID NO: 56) or Figures 25A-25D (SEQ ID NO: 55), and which GABA<sub>B</sub>R2 polypeptide has an amino acid sequence (a) identical to the amino acid sequence shown in Figures 4A-4D (SEQ ID NO: 4) or Figures 23A-23D (SEQ ID NO: 47), or (b) encoded by a nucleic acid sequence identical to the receptor-encoding nucleic acid sequence contained in plasmid pEXJT3T7-hGABAB2

G<sup>5</sup>  
cont

(ATCC Accession No. 203515) or in plasmid BO-55 (ATCC Accession No. 209104).--

-213. (Three Times Amended) A process for determining whether a chemical compound activates a mammalian GABA<sub>B</sub>R1/R2 receptor, which comprises contacting cells producing a second messenger response and expressing on their cell surface the GABA<sub>B</sub>R1/R2 receptor, wherein such cells prior to being transfected with such nucleic acid do not express the GABA<sub>B</sub>R1/R2 receptor, with the chemical compound under conditions suitable for activation of the GABA<sub>B</sub>R1/R2 receptor, and measuring the second messenger response in the presence and in the absence of the chemical compound, a change in the second messenger response in the presence of the chemical compound indicating that the compound activates the GABA<sub>B</sub>R1/R2 receptor, wherein the mammalian GABA<sub>B</sub>R1/R2 receptor comprises a GABA<sub>B</sub>R1 polypeptide and a GABA<sub>B</sub>R2 polypeptide, which GABA<sub>B</sub>R1 polypeptide has an amino acid sequence identical to the amino acid sequence shown in Figures 24A-24D (SEQ ID NO: 56) or Figures 25A-25D (SEQ ID NO: 55), and which GABA<sub>B</sub>R2 polypeptide has an amino acid sequence (a) identical to the amino acid sequence shown in Figures 4A-4D (SEQ ID NO: 4) or Figures 23A-23D (SEQ ID NO: 47), or (b) encoded by a nucleic acid sequence identical to the receptor-encoding nucleic acid sequence contained in plasmid pEXJT3T7-hGABAB2 (ATCC Accession No. 203515) or in plasmid BO-55 (ATCC Accession No. 209104).--

G<sup>6</sup>

-224. (Four Times Amended) A method of screening a plurality of chemical compounds to determine whether any

G<sup>7</sup>

compound within such plurality of chemical compounds activates the GABA<sub>B</sub>R1/R2 receptor, wherein the mammalian GABA<sub>B</sub>R1/R2 receptor comprises a GABA<sub>B</sub>R1 polypeptide and a GABA<sub>B</sub>R2 polypeptide, which GABA<sub>B</sub>R1 polypeptide has an amino acid sequence identical to the amino acid sequence shown in Figures 24A-24D (SEQ ID NO: 56) or Figures 25A-25D (SEQ ID NO: 55), and which GABA<sub>B</sub>R2 polypeptide has an amino acid sequence (a) identical to the amino acid sequence shown in Figures 4A-4D (SEQ ID NO: 4) or Figures 23A-23D (SEQ ID NO: 47), or (b) encoded by a nucleic acid sequence identical to the receptor-encoding nucleic acid sequence contained in plasmid pEXJT3T7-hGABAB2 (ATCC Accession No. 203515) or in plasmid BO-55 (ATCC Accession No. 209104) which comprises:

- G<sup>7</sup>  
Cont.
- (a) contacting cells containing nucleic acid encoding, and expressing on their cell surface, the GABA<sub>B</sub>R1/R2 receptor, wherein such cells prior to being transfected with such nucleic acid do not express the GABA<sub>B</sub>R1/R2 receptor, with the plurality of compounds, under conditions permitting activation of the GABA<sub>B</sub>R1/R2 receptor;
  - (b) determining whether the activity of the GABA<sub>B</sub>R1/R2 receptor is increased in the presence of the compounds, and if it is increased;
  - (c) separately determining whether the activation of the GABA<sub>B</sub>R1/R2 receptor is increased by each compound included in the plurality of compounds, so as to thereby determine whether any compound or compounds present in such plurality of compounds activates the

G2

GABA<sub>B</sub>R1/R2 receptor. --

G8

-230. (Amended) The method of claim 250, wherein the non-neuronal cell is a COS-7 cell, a 293 human embryonic kidney cell, a LM(tk-) cell or an NIH-3T3 cell --

G9

-231. (Three Times Amended) A process for determining whether a chemical compound is an agonist of a mammalian GABA<sub>B</sub>R1/R2 receptor, which comprises preparing a membrane fraction from cells which comprise nucleic acid encoding, and expressing on their cell surface, the GABA<sub>B</sub>R1/R2 receptor, wherein such cells prior to being transfected with such nucleic acid do not express the GABA<sub>B</sub>R1/R2 receptor, separately contacting the membrane fraction with both the chemical compound and GTPγS, and with only GTPγS, under conditions permitting the activation of the GABA<sub>B</sub>R1/R2 receptor, and detecting GTPγS binding to the membrane fraction, an increase in GTPγS binding in the presence of the compound indicating that the chemical compound activates the GABA<sub>B</sub>R1/R2 receptor, wherein the mammalian GABA<sub>B</sub>R1/R2 receptor comprises a GABA<sub>B</sub>R1 polypeptide and a GABA<sub>B</sub>R2 polypeptide, which GABA<sub>B</sub>R1 polypeptide has an amino acid sequence identical to the amino acid sequence shown in Figures 24A-24D (SEQ ID NO: 56) or Figures 25A-25D (SEQ ID NO: 55), and which GABA<sub>B</sub>R2 polypeptide has an amino acid sequence (a) identical to the amino acid sequence shown in Figures 4A-4D (SEQ ID NO: 4) or Figures 23A-23D (SEQ ID NO: 47), or (b) encoded by a nucleic acid sequence identical to the receptor-encoding nucleic acid sequence contained in plasmid pEXJT3T7-hGABAB2 (ATCC Accession No. 203515) or in plasmid BO-55 (ATCC